

RESEARCH

Hip Knee J. Vol. 3, No. 2, 2022, pp. 64-70

p-ISSN: 2723-7818

e-ISSN: 2723-7826

<http://dx.doi.org/10.46355/hipknee.v3i2.130>

CORRELATION REALIGNMENT MECHANICAL TIBIOFEMORAL ANGLE WITH CLINICAL OUTCOME ON PATIENTS KNEE OSTEOARTHRITIS VALGUS DEFORMITY PERFORMED TOTAL KNEE ARTHROPLASTY AT PROF. DR. R. SOEHARSO ORTHOPAEDIC HOSPITAL SURAKARTA (JANUARY 2019 - DECEMBER 2019)

Umar Kharisma Islami¹, Fanny Indra Warman¹, Pamudji Utomo¹, Iwan Budiwan Anwar¹, Tangkas SMHS Sibarani¹

¹Department Orthopaedic and Traumatology, Universitas Sebelas Maret, Prof.Dr.R.Soeharso, Orthopaedic Hospital, Surakarta

ABSTRACT

Background: Knee osteoarthritis (OA) are the most important because of the high prevalence of pain and disability in older adults. Angular deformities around the knee joint necessitate special consideration to restore normal alignment during TKA. In the region of 10–15% of patients requiring a primary TKA present with a valgus deformity (VD), the accurate correction of which still poses a challenge. Excessive preoperative malalignment predisposes to a greater risk of failure compared to well-aligned knees. In Ranawat grade-III the axis deviation is more than 20°. All the medial stabilizing elements are typically not functional so a constrained implant usually is required. The aim of this research to savvy correlation realignment mechanical tibiofemoral angle with clinical outcome on patients knee osteoarthritis valgus deformity performed TKA with non-constrained implant.

Methods: This research is an analytics observational with *cross sectional* that group patients knee osteoarthritis valgus devormity underwent total knee arthroplasty and performed correlation realignment tibiofemoral angle as well as clinical outcome. This research with total sampling method on patients knee osteoarthritis valgus deformity Kellgren-Lawrence grade III–IV, have radiological knee x-ray and lower extremity scanogram underwent TKA one side by orthopaedic surgeon sub adult reconstruction at Prof. Dr. R. Soeharso orthopaedic hospital Surakarta on periode January 2019 until Desember 2019. Evaluation has performed to pain scale used Visual Analog Scale (VAS) score, clinical outcome with Knee Society Score (KSS), realignment mechanical tibiofemoral angle before and after TKA. Data were analyzed using t-test and correlation test with SPSS version 16.

Results: Analyzed 28 (male 6, female 22) patients with a mean age 65,75 years, which divided into three group Ranawat grade I was 14 patients, grade II was 9 patients and III was 5 patients. They were evaluated by VAS score, KSS score, mechanical tibiofemoral angle before and after TKA. Evaluated for VAS score, KSS objective and functional, mechanical tibiofemoral angle have significant result ($P < 0,05$). There was also significantly correlation between improvement mechanical tibiofemoral angle with KSS knee score based value ($P < 0,05$).

Conclusion: In this study we found that TKA with non-constrained implant on knee osteoarthritis valgus deformity provides significant result in reducing pain scale and increased clinical outcome.

Keywords: *Knee osteoarthritis valgus deformity, Total Knee Arthroplasty, Knee Society Score, VAS Score, mechanical tibiofemoral angle*

Article history:

Submission : May 10th, 2022
Revision : June 10th, 2022
Accepted : June 23th, 2022

Corresponding Author:

Pamudji Utomo, MD, PhD
Prof. Dr. R. Soeharso Orthopaedic Hospital, Jl. Jenderal
Ahmad Yani, Pabelan, Sukoharjo, 57162
email: pamudjiutomo@gmail.com

Introductions

Osteoarthritis (OA) is by far the most common form of joint disease throughout the world. It is strongly associated with age, and extremely common in older people; some studies estimate that over 80% of people over 55 years of age have osteoarthritis of at least one joint. Hip and knee OA are the most important because of the high prevalence of pain and disability that they cause in older adults, and the massive healthcare resource input that results from this, particularly in terms of the provision of joint replacements.⁽¹⁾

Angular deformities around the knee joint necessitate special consideration to restore normal alignment during total knee arthroplasty (TKA). In the region of 10–15% of patients requiring a primary TKA present with a valgus deformity (VD), the accurate correction of which still poses a challenge.⁽²⁾⁽³⁾ Excessive preoperative malalignment predisposes to a greater risk of failure compared to well-aligned knees.⁽⁴⁾

Approximately 10% of patients requiring total knee arthroplasty have a valgus deformity (defined as an anatomic valgus of $>10^\circ$). Correction of the valgus deformity has posed technical challenges and has produced variable clinical results in terms of correction of the deformity, instability, and the overall results.⁽⁵⁾

The valgus deformity in osteoarthritis knee is a continuing process developed by bone tissue remodeling and soft tissue contraction/elongation. Bone tissue alterations consist of lateral cartilage erosion, lateral condylar hypoplasia, and metaphyseal femur and tibial plateau remodeling. Soft tissue alterations are characterized by tightening of lateral structures such as lateral collateral ligament (LCL), posterolateral capsule (PLC), popliteus tendon (POP), hamstring tendons, the lateral head of the gastrocnemius (LHG) and iliotibial band (ITB).^(5,6)

In Ranawat grade III the axis deviation is greater than 20° . All stabilizing elements, especially the medial section does not function, implant constrained is needed.^(2,3)

The purpose of this research to savvy correlation realignment mechanical tibiofemoral angle with clinical outcome on patients knee osteoarthritis valgus deformity performed TKA with non-constrained implant. Based on hypothesis there is difference between the VAS (Visual Analog Score) patients preoperative and postoperative TKA, difference between the MTFA (mechanical tibiofemoral angle) patients preoperative and postoperative TKA, correlation between improvement MTFA with improvement KSS preoperative and postoperative TKA.

Methods

This research is an analytics observational with cross sectional that group patients knee osteoarthritis valgus devormity underwent total knee arthroplasty and performed correlation realignment tibiofemoral angle as well as clinical outcome with KSS. Data were collected from the patient data bank in the sub adult reconstruction departement which is synchronized with the patient's medical record data at Prof. Dr. R. Soeharso Surakarta Orthopedic Hospital

This research with total sampling method with criteria inclusion patients knee osteoarthritis valgus deformity Kellgren-Lawrence grade III–IV, have radiological knee x-ray and lower extremity scanogram underwent TKA one side by orthopaedic surgeon sub adult reconstruction at Prof. Dr. R. Soeharso orthopaedic hospital Surakarta on periode January 2019 until Desember 2019. All surgeries were performed by 4 senior orthopaedic surgeons who were adult reconstruction specialists.

Exclusion criteria were performed, patients with total revision of knee arthroplasty, patients who had deformity abnormalities in the lower extremities, patients died, patients refused to be used as study samples.

Evaluation has performed to pain scale used Visual Analog Scale (VAS) score, clinical outcome with Knee Society Score (KSS), realignment mechanical tibiofemoral angle before and after TKA. Data analysis of the research using

normalitas test with kolmogorov-smirnov test. Data were analyzed using t-test and correlation test with SPSS version 16.

Result

All patients with osteoarthritis of the knee valgus deformity who underwent total knee arthroplasty surgery according to exclusion criteria, there were 4 patients to follow-up loss, 2 patients did not have a scanogram data (1 patients of inadequate scanogram and 1 patient had an infection and did not stand properly). Total inclusion criteria was 28 patients.

This study population consist of 22 women (78.6%) and 6 man (21.4%). The mean age patient was 65 year (range 46 – 77 years), whereas the mean BMI 26.07 kg/m² (range, 20.03 – 36.68 kg/m²). Side operation consist of right side 21 (75%) and left side 7 (25%). Mean data of postoperative evaluation was 4.7 months (2 – 10 months).(Tabel 1).

Ranawat grade distribution consist of grade I was 14 (50%), grade II was 9 (32.1%) and grade III was 5 (17.9%). The mean pre operative knee joint ROM was 105.18 ° (range, 90° - 115°), post operative knee joint ROM was 106.25 ° (range, 70° - 130°). Based on the Ranawat Grade, in Ranawat Grade I, post operative knee joint ROM was 108.9° (range, 80° - 130°); in Ranawat Grade II, post operative knee joint ROM was 96.5° (range, 70° - 120°); whereas in Ranawat Grade III, post operative knee joint ROM was 118° (range, 115° - 120°).

Tabel 1. Preoperative demographic of the patient

Sex (male/female)	6/22
Age (y)	65 (46 – 77)
Body mass index (kg/mm2)	26.07 (20.03 – 36.68)
Normal	15 (50%)
Mild obese	6 (21%)
Obese	8 (29%)
Range of motion	105.18° (90° - 115°)
MTFA	11.35° (1° - 27°)
Operation side (right/left)	21/7
Follow up (months)	4.7 (2 – 10)
Ranawat grade	28
Grade I	14 (50%)
Grade II	9 (32.1%)
Grade III	5 (17.9%)

Clinical and Radiological Outcome

The mean VAS score pre operative was 6.39 (range, 5 – 7), whears the mean post operative was 1.42 (range, 1-3). KSS (Knee Society Score) consist of KSS knee score pre operative was 28 poor (100%), and post operative with criteria 1 fair (3.5%), 8 good (28.5%), 19 excellent (67.8%). KSS function score pre operative was 28 poor (100%), and post operative with criteria 1 poor (3,5%), 2 fair (7.1%), 20 good (71.4), 5 excellent (17.8%).

The VAS score, MTFA, and KSS post operative score were significantly improved at evaluation compared pre operative baseline value (p < 0,05). However for ROM knee joint post operative has no significantly improved compared post operative baseline value (p < 0,05) (Tabel 2).

Tabel 2. Comparison of mean pre operative and post operative clinical and radiological result

	Pre operative	Post operative	P-value
VAS score	6.39	1.42	P < 0.05
ROM knee joint	104.46	106.25	P > 0.05
MTFA	11.35	-0.87	P < 0.05
KSS score			
KSS knee score	44.17	81.14	P < 0.05
KKS function score	31.78	70.17	P < 0.05

There is no significant comparison between Ranawat grade 1 and 2/3 with KSS knee score as well as KSS function score baseline value (p > 0,05).

There is significantly correlation between improvement MTFA with improvement KSS knee score osteoarthritis valgus deformity preoperative and postoperative baseline value (p < 0,05). (Figure 1). There is no correlation between improvement MTFA with improvement KSS function score osteoarthritis valgus deformity preoperative and postoperative baseline value (p > 0,05). Data after operation experienced under correction of MTFA was 14 patient (50%) with mean 2.78° valgus, neutral correction was 1

patient (3.5%) and over correction was 13 (46.5%) with mean 3.61° varus. (Tabel 3). There is result post operative correction MTFA based grade Ranawat. Valgus correction Ranawat grade I (5), II (5), III (4). Varus correction Ranawat grade I (8), II (4), III (1) (Tabel 4). There is no difference between realignment of knee osteoarthritis valgus deformity that become *valgus* and *varus* with KSS post operative baseline value ($p > 0,05$).




Figure 1. Measurement of MTFA pre operative (22° valgus) and post operative (2°) realignment correction of knee OA valgus deformity on lower extremity scanogram.

Tabel 3. Result post correction MTFA

Preoperative valgus	Post operative		
	Valgus	Neutral	Varus
28 (100%)	14 (50%)	1 (3.5%)	13 (46.5%)

Tabel 4. Result post operative correction MTFA based grade Ranawat

Pre operative Ranawat	Post operative correction	
	Valgus	Varus
Grade I	5	8
Grade II	5	4
Grade III	4	1

Discussion

In the study for the average age of patients around 65.75 years. Based on research conducted by Van Schoor et al, found that people aged 65-85 years have a 1.5 times higher risk of developing OA than other ages.⁽⁷⁾ De Zwart, et al also found the risk of a decrease in muscle strength in old age as predictive factors for the occurrence of OA.⁽⁸⁾

From the data it was found that normal body mass index 14 (50%), mild obesity 6 (21%) severe obesity 8 (29%). Based on research conducted by Kumar et al, it was found that the obese population has a tendency to experience OA by 33% compared to others. This percentage will increase according to the increase in BMI.⁽⁹⁾ Burn et al in his study also found that increasing BMI will increase a person to risk of developing knee OA.⁽¹⁰⁾

Based on the data obtained from 28 patients, there were 22 female sex with a percentage of 78.6% and 6 male sex with a percentage of 21.4%. This result is in accordance with research conducted by Kumar et al. which states that statistically more women experience OA than men due to the tendency of women to be obese compared to men.⁽⁹⁾ This conformity was also found in Pontoh et al, study which found a ratio of women compared to men who experienced OA as much as 3: 1.⁽¹¹⁾ Based on research by Jung et al, found that postmenopausal women have decreased estrogen levels which causes accelerated damage from the knee cartilage thereby increasing the risk of OA. Low estradiol content in postmenopausal women will increase the pain threshold in OA.⁽¹²⁾

In the postoperative phase, all 28 patients were in the mild pain category (VAS: 1-3) with a percentage of 100%. In Ranawat grade 1, 2 and 3 all gave the same VAS score, that is the mild pain category. This is consistent with the research conducted by Ramlall et al. In which he reported pain scores in TKR pre operative patients had a significant result compared to after surgery. Ramlall et al found an average pain score in patients before surgery of 7 out of 10, and an average pain score after surgery of 3 out of 10.⁽¹³⁾

One of the main indications of TKA is persistent pain that is often found in end-stage OA patients which cannot be treated with non-invasive methods.^(14,15) In this study there was a significant reduction in pain.

The comparative analysis results of this study between the preoperative range of motion of the knee joint 104.45° and 106.25° postoperatively showed a *P* value (significance) of 0.737, which means $P > 0.05$. This showed that the comparative test between the 2 groups was not significant. In the comparison of patients with osteoarthritis Ranawat grade 1, 2 and 3 for improvement knee joint range of motion, that there was no significant difference between the groups. According to the study of Redulescu et al, there was an increase in range of motion post knee osteoarthritis surgery with severe valgus deformity obtained an average result of preoperative 71° and postoperative 95° .⁽¹⁶⁾ This result is different from the study of Redulescu et al. It is possible that the factor in the existence of patients who are less cooperative in undergoing physiotherapy. That is can caused less optimal knee range of motion. In patients who underwent total knee arthroplasty, surgery has been evaluated to obtain optimal knee ROM.

The results of the different tests between the KSS knee score and the KSS function score preoperatively and postoperatively showed a *P*-value (significance) of 0,000, which means $P < 0.05$. This showed that the postoperative KSS experienced a significant improvement compared to the preoperative KSS. This is in accordance with research Redulescu et al who also found that knee osteoarthritis surgery with severe valgus deformity, KSS experienced an improved from a preoperative average of 21.3° to an average of 80.7° postoperatively.⁽¹⁶⁾ That means it could increased clinical outcomes in patients with knee osteoarthritis with valgus deformity who have performed total knee arthroplasty with non-constrained implants.

In this study, the test results different between the mechanical tibiofemoral angle (MTFA) on average preoperatively by 11.35°

valgus and postoperative average of 0.17° varus. The *P* value (significance) is 0,00, which means $P < 0.05$. This showed that MTFA experienced a significant change between pre and post operative. Mechanically aligned TKA's aim to have a neutral coronal plane alignment and a tibial cut that is orthogonal to the mechanical axis. In a valgus knee, both femoral and tibial pathology may need to be addressed with respective bone cuts to achieve this goal.⁽³⁾ In this study the total knee arthroplasty is underwent cutting both the tibia and the femoral is using templating, it is expected to restore the alignment of the neutral coronal plane.

In this study the relationship between improvement MTFA with improvement of the KSS Knee Score shows the *P* value (significance) is 0,000, which means $P < 0.05$, so it can be concluded that MTFA improvement is significantly related to improvement in the KSS Knee Score. Choong 2009 mentions that patients who achieved alignment in the mechanical angle 3° tibiofemoral axis increased the KSS. Orthopedic experts believe that good alignment (in 3° mechanical axis) TKA results in improved outcomes.⁽⁹⁾ One goal for TKA is to achieve good results in alignment of the femur, tibia and patella components by restored the patient's lower extremities to neutral. Proper alignment of the knee joint is considered as one of the most influential factors in determining the long-term outcome after TKA.⁽¹⁰⁾ In this study the results of post-correction MTFA with mean valgus 2.78° and varus 3.61° .

Based on this study clinical outcome evaluation did not differ during the short-term evaluation, it needed a long-term follow-up evaluation to assess the clinical outcome. In this study the actioned of TKA with non-constrained implants in the valgus deformity are quite reliable with the requirement that the realignment correction performed can be achieved.

Complication

Complications were found in this study. One patient experienced infection after operation

and did not to stand properly so excluded to the study. Another complication was lack of knee joint ROM after surgery. There were 4 patients post operative with less than 90° knee joint ROM. Erkan et al mentioned that stiffness in the knee joint was a postoperative complication that occurred in 4-16% of patients with TKA and was a significant cause of morbidity due to pain and limited function. Preoperative flexion range is limited, then the stiff knee is more likely to occur in patients who will undergo TKA. ^(17,18) Risk factors for stiffness after TKA can be divided into 3 groups: preoperative factors, intraoperative factors and post operative factors. ⁽¹⁹⁾ In these patients it is possible due to postoperative factors such as inadequate rehabilitation or physiotherapy with low patient motivation.

Limitation

Some limitation of this study are the limited number of patient samples, which is less than 30 patients. Evaluation of postoperative patient follow-up is quite short under 1 year, research is needed to continue with long-term evaluation. In addition, there is haven't detailed data about the surgical technical procedures that have been performed.

Conclusions

Based on this study total knee arthroplasty with non-constrained implant on knee osteoarthritis valgus deformity provides significant result in reducing pain scale and increased clinical outcome. TKA with non-constrained implants in the valgus deformity are quite reliable with the requirement that the realignment correction performed can be achieved.

References

1. Blom, A. Dieppe, P. Osteoarthritis in Apley & Solomon's System of Orthopaedics and Trauma. 10th ed. CRC Press. New York, US. 2018.
2. Dimitrios, N. George, S. Ioannis, M. Total Knee Arthroplasty in Valgus Deformity. *Chapter 4. IntechOpen*. 2018.
3. Lange J, Haas SB. Management factorials in primary total knee arthroplasty correcting severe valgus deformity taking out the knock. *Bone Jt J*. 2017;99B(1):60–4.
4. Rossi R, Rosso F, Cottino U, Dettoni F, Bonasia DE, Bruzzone M. Total knee arthroplasty in the valgus knee. *Int Orthop*. 2014;38(2):273–83.
5. Ranawat AS, Ranawat CS, Elkus M, Rasquinha VJ, Rossi R, Babhulkar S. Total Knee Arthroplasty for Severe Valgus Deformity Surgical Technique. *J Bone Jt Surgery, Inc Jbjs*. 2004;86:2671–6.
6. Utomo DN, Mahyudin F, Yanuar A, Widhiyanto L, Hernugrahanto KD. Correction of severe valgus deformity of knee osteoarthritis with non-constrained total knee arthroplasty implant: A case report. *Int J Surg Case Rep*. 2018;53:218–22.
7. van Schoor NM, Dennison E, Castell M V., Cooper C, Edwards MH, Maggi S, et al. Clinical osteoarthritis of the hip and knee and fall risk: The role of low physical functioning and pain medication. *Semin Arthritis Rheum*. 2020.
8. de Zwart AH, van der Leeden M, Roorda LD, van der Esch M, Twisk JWR, Lems WF, et al. Course and predictors of upper leg muscle strength over 48 months in subjects with knee osteoarthritis: Data from the osteoarthritis initiative. *Osteoarthr Cartil Open [Internet]*. 2020;(October 2019):100038.
9. Kumar H, Pal CP, Sharma YK, Kumar S, Uppal A. Epidemiology of knee osteoarthritis using Kellgren and Lawrence scale in Indian population. *J Clin Orthop Trauma [Internet]*. 2020;11(xxxx):S125–9.
10. Burn E, Murray DW, Hawker GA, Pinedo-Villanueva R, Prieto-Alhambra D. Lifetime risk of knee and hip replacement following a GP diagnosis of osteoarthritis: a real-world cohort study. *Osteoarthr Cartil*.

- 2019;27(11):1627–35.
11. Pontoh LA, Putra A, Dologo IH, Efar TS. Characteristics of coexisting patellofemoral joint osteoarthritis and tibiofemoral joint osteoarthritis in an Indonesian population: A cross-sectional study at a tertiary teaching hospital. *Int J Surg Open*. 2019;21:64–7.
 12. Jung JH, Bang CH, Song GG, Kim C, Kim JH, Choi SJ. Knee osteoarthritis and menopausal hormone therapy in postmenopausal women: A nationwide cross-sectional study. *Menopause*. 2019;26(6):598–602.
 13. Ramlall Y, Andrion JJD, Cameron HU, Sawhney M. Examining pain before and after primary total knee replacement (TKR): A retrospective chart review. *Int J Orthop Trauma Nurs* [Internet]. 2019;34(April):43–7.
 14. Drosos GI, Triantafilidou T, Ververidis A, Agelopoulou C, Vogiatzaki T, Kazakos K. Persistent post-surgical pain and neuropathic pain after total knee replacement. *World J Orthop*. 2015;6(7):528–36.
 15. Price AJ, Alvand A, Troelsen A, Katz JN, Hooper G, Gray A, et al. *Knee Replacement*. *Lancet Lond Engl*. 03 2018;392(10158):1672–82.
 16. Radulescu R, Badila A, Japie I, Ciobanu T, Manolescu R. Primary total knee arthroplasty in severe valgus knee. *J Med Life*. 2013;6(4):395–8.
 17. Erkan S, Yercan HS, Okcu G, Ozalp RT. Factors causing stiff knee after total knee arthroplasty. *Eklemler Hastalik Cerrahisi* 2011;22:16–21.
 18. Ghani H, Maffulli N, Khanduja V. Management of stiffness following total knee arthroplasty: A systematic review. *Knee*. 2012;19(6):751–9.
 19. Rodríguez-Merchán EC. The stiff total knee arthroplasty: Causes, treatment modalities and results. *EFORT OpenRev*. 2019;4(10):602–10.